arXiv:funct-an/9211005v1 25 Nov 1992

A Short Proof that $M_n(A)$ is local if A is local and Fréchet

Larry B. Schweitzer

Current address: Department of Mathematics, University of Victoria
Victoria, B.C. Canada V8W 3P4
Email Address: lschweit@sol.uvic.ca

Abstract

We give a short and very general proof of the fact that the property of a dense Fréchet subalgebra of a Banach algebra being local, or closed under the holomorphic functional calculus in the Banach algebra, is preserved by tensoring with the $n \times n$ matrix algebra of the complex numbers.

Introduction

We say that a dense subalgebra A of a Banach algebra B, having the same unit as B, is local [**2**], or closed under the holomorphic functional calculus in B if for each $a \in A$ and each function f holomorphic in a neighborhood of the spectrum of a in B, the element $f(a)$ of B lies in A. If A is local in B, then the inclusion map $\iota\colon A \hookrightarrow B$ induces an isomorphism of K-theories $\iota_*\colon K_*(A) \cong K_*(B)$ [**5, VI.3**], [**3, Theorem A.2.1**]. This is one indication of the importance of local dense subalgebras.

In §1, we introduce the notion of spectral invariance, and show that, if A is a Fréchet algebra under some possibly stronger topology than that of B, then A is spectral invariant in B iff A is local in B. We then show that A is spectral invaraint in B iff every simple A-module (or algebraically irreducible representation of A) is contained in a B-module, and use this to give a simple proof that the $n \times n$ matrices $M_n(A)$ are spectral invariant in $M_n(B)$ iff A is spectral invariant in B. This result is useful in K-theory [**10, Corollary 7.9**]. We conclude with several interesting examples. This paper is a portion of my Ph.D. thesis, written at the University of California at Berkeley, under the supervision of Marc A. Rieffel.

§1 An Equivalent Condition for Spectral Invariance.

We show in this section that a dense subalgebra A of a topological Q-algebra B is spectral invariant in B if and only if all simple A-modules are contained in B-modules.

Definition 1.1. By a topological algebra, we mean a topological vector space over a topological ring k with unit, with a k-algebra structure for which multiplication is separately continuous. Let B be a topological algebra over a topological ring k with unit. Let A be a subalgebra. If B has no unit let $\tilde{B}$ be B with unit adjoined, and let $\tilde{A}$ be A with the same unit adjoined. If B has a unit, and A has the same unit, we let $\tilde{A} = A$, $\tilde{B} = B$. If B has a unit 1_B and A has a different unit, we let $\tilde{B} = B$ and $\tilde{A}$ be the subalgebra of B generated by A and 1_B. Note that in any case $\tilde{B}$ is then a topological algebra, and $\tilde{A}$ a subalgebra. If A is dense in B, then $\tilde{A}$ is dense in $\tilde{B}$ and B has the same unit as A if A is unital. We often make the additional assumption that the group $\mathrm{inv}\tilde{B}$ of invertible elements in $\tilde{B}$ is open. If $\mathrm{inv}\tilde{B}$ is open, we call B a *Q-algebra* (see [**9**]). If, in addition, inversion in $\tilde{B}$ is continuous, we say that B is a *good* topological algebra (see [**3**]).

We say that A is a *spectral invariant (SI)* subalgebra of B if the invertible elements of $\tilde{A}$ are precisely those elements of $\tilde{A}$ which are invertible in $\tilde{B}$. Note that A is SI in B if and only if for every $a \in A$, the spectrum of a is the same in $\tilde{A}$ and in $\tilde{B}$.

By a Fréchet algebra, we shall mean a Fréchet space with an algebra structure for which multiplication is jointly continuous [**13**]. We say that a Fréchet algebra is m-convex if its topology is given by a family of submultiplicative seminorms. An m-convex Fréchet algebra has a holomorphic functional calculus [**10, Lemma 1.3**]. If B is an m-convex Fréchet algebra and A is a subalgebra, we say that A *is closed under the holomorphic functional calulus in* B (or *local in* B [**2**]) if and only if for every $a \in \tilde{A}$ and f holomorphic in a neighborhood of $spec_{\tilde{B}}(a)$, the element $f(a)$ of $\tilde{B}$ lies in $\tilde{A}$.

We say that A is a *Fréchet subalgebra* of a topological algebra B if A is a Fréchet algebra, and there exists an injective and continuous algebra homomorphism (called the inclusion map) $\iota\colon A \hookrightarrow B$.

Lemma 1.2. *Let B be an m-convex Fréchet Q-algebra and A a Fréchet subalgebra. Then A is closed under the holomorphic functional calculus in B iff A is spectral invariant in B.*

Proof: For $\Rightarrow$, note that if $a \in \mathrm{inv}\tilde{B} \cap \tilde{A}$, then $f(\lambda) = 1/\lambda$ is holomorphic in a neighborhood of the closed set $spec_{\tilde{B}}(a)$ (compare [**2, Proposition 3.1.3**]). (We use $\mathrm{inv}\tilde{B}$ open to see that $spec_{\tilde{B}}(a)$ is closed.)

For $\Leftarrow$ first note that $\mathrm{inv}\tilde{A} = \mathrm{inv}\tilde{B} \cap \tilde{A}$ is open since $\mathrm{inv}\tilde{B}$ is open and the inclusion map from A to B is continuous. It follows that A has holomorphic functional calculus [**8**] [**13**] which agrees with the holomorphic functional calculus on B. This fact is well known, but I have not been able to find it stated in the literature, so I will prove it here for convenience. Since $\mathrm{inv}\tilde{A}$ is open and $\tilde{A}$ is a Fréchet algebra, the inverse map $x \mapsto x^{-1}$ in $\mathrm{inv}\tilde{A}$ is continuous [**13, Chapter 7, Proposition 2**]. If f is holomorphic in a neighborhood N of $spec_{\tilde{B}}(a)$, then $\lambda \mapsto f(\lambda)(\lambda - a)^{-1}$ is a well defined and continuous map from any closed curve Γ contained in $N - spec_{\tilde{B}}(a)$. From this one checks that the integral expression

$$f(a) = \frac{1}{2\pi\imath} \int_{\Gamma} f(\lambda)(\lambda - a)^{-1} d\lambda$$

has its partial Riemann sums converging in $\tilde{A}$. Thus $f(a) \in \tilde{A}$.**Q.E.D.**

We shall often call a simple module an irreducible representation. Irreducible will mean algebraically irreducible. We shall use the terminology module-representation and submodule-subrepresentation interchangably.

To motivate Theorem 1.4, we consider the situation where the algebra B and its dense subalgebra A are commutative and unital Banach algebras. Then all simple A-modules are in fact homomorphisms of A to $\mathbf{C}$. If a is not invertible in A, one can easily construct a homomorphism from A to $\mathbf{C}$ that has a in its kernel. If this homomorphism extends to a homomorphism from B to $\mathbf{C}$, clearly a cannot be invertible in B (so if all homomorphisms of A extend to ones of B, A must be spectral invariant in B).

Conversely, let h be a homomorphism of A to $\mathbf{C}$ with no extension to a homomorphism of B. Assume that the kernel M of h is not dense in B. Then the closure $\overline{M}$ is a proper ideal in B. By Zorn's lemma, $\overline{M}$ is contained in a maximal closed ideal N in B. The quotient map

$$B \to B/N \cong \mathbf{C} \qquad (\mathrm{Gelfand-MazurTheorem})$$

gives an extension of the homomorphism h (it extends h since $N \cap A = M$ by the maximality of M as an ideal in A). So since the homomorphism h has no extension, our assumption that M is not dense in B has led us to a contradiction. So M must be dense in B. Since B

is a Q-algebra, M must contain an invertible element of B which is contained in A. Since M is a proper ideal of A, this element is not invertible in A, and thus A is not spectral invariant in B. So A is SI in B iff every simple A-module is contained in a B-module. We will generalize these observations in Theorem 1.4.

In some literature ([**3**], [**12**]), the following more general setting is considered.

Definition 1.3. Let B be a topological algebra and let $\varphi\colon A \to B$ be an algebra homomorphism. We define $\tilde{A}$ and $\tilde{B}$ as in Definition 1.1. We say that A is *spectral invariant* in B if for all $a \in \tilde{A}$, a is invertible in $\tilde{A}$ whenever $\varphi(a)$ is invertible in $\tilde{B}$. We say that an A-module V is *contained in* a B-module W if $V \subseteq W$, V factors to a $\varphi(A)$-module, and the action of B on W is consistent with the action of $\varphi(A)$ on V.

We prove the following theorem in this setting. It is similar to [**7, Theorem 2.4.16**].

Theorem 1.4. *Let B be a Q-algebra, and let $\varphi\colon A \to B$ be an algebra homomorphism with dense image. The following are equivalent:*

(i) A is spectral invariant in B.

(ii) For every maximal left ideal N in $\tilde{A}$, we have $\varphi^{-1}(\overline{\varphi(N)}^{\tilde{B}}) = N$.

(iii) Every simple A-module V is contained in a B-module W. If so desired, W can be taken to be a simple B-module.

The proof of $(iii) \Rightarrow (i)$ does not require $\mathrm{inv}\tilde{B}$ *open or $\varphi(A)$ dense.*

Also, the B-module W constructed in our proof (of $(ii) \Rightarrow (iii)$) inherits a natural topology, making it a continuous B-module for which the action of B is continuous (more precisely, the map $(b, w) \mapsto bw$ from $B \times W$ to W is separately continuous). In this topology, V is dense in W.

Proof:

$(i) \Rightarrow (ii)$. Assume A is SI in B. Let N be a maximal left ideal in $\tilde{A}$. Extend φ in the natural way to a map $\varphi\colon \tilde{A} \to \tilde{B}$, again having dense image. Then $N \cap \mathrm{inv}\tilde{A} = \emptyset$ so $\varphi(N) \cap \mathrm{inv}\tilde{B} = \emptyset$ by spectral invariance. Since $\mathrm{inv}\tilde{B}$ is open in $\tilde{B}$, the closure $\overline{\varphi(N)}$ of $\varphi(N)$ in $\tilde{B}$ is a proper subset of $\tilde{B}$, and is also a left ideal in $\tilde{B}$. Since $\varphi(\tilde{A})$ is dense in

$\tilde{B}$, $\varphi^{-1}(\overline{\varphi(N)}) \neq \tilde{A}$. Thus $\varphi^{-1}(\overline{\varphi(N)})$ is a proper left ideal in $\tilde{A}$. Since this ideal contains N and N is maximal, we have $\varphi^{-1}(\overline{\varphi(N)}) = N$. (Note that any maximal left ideal of $\tilde{A}$ therefore contains the kernel of φ.)

(ii) $\Rightarrow$ *(iii)*. Let V be a simple A-module. Extend V to a $\tilde{A}$-module in the natural way. Let $v \in V$, $v \neq 0$. Let N be the kernel of the map $\tilde{A} \to V$, $a \mapsto av$. This map is onto by simplicity, so $V \cong \tilde{A}/N$ as A modules. Also by simplicity, N is a maximal left ideal, so by *(ii)* we have $\varphi^{-1}(\overline{\varphi(N)}) = N$. Set $M = \overline{\varphi(N)}$. Then the map $\varphi\colon \tilde{A} \to \tilde{B}$ gives the inclusion

$$V \cong \tilde{A}/N \subseteq \tilde{B}/M \qquad (*)$$

where $\tilde{B}/M$ is a B-module. Take $W = \tilde{B}/M$. This proves *(ii)* $\Rightarrow$ *(iii)*.

Clearly M is a closed ideal in $\tilde{B}$ so W inherits a natural topology in which V is dense. If we replace M with a maximal ideal in $\tilde{B}$ containing $\overline{\varphi(N)}$, we still have $\varphi^{-1}(M) = N$ and so $(*)$ continues to hold. Also M must be closed, so $W = \tilde{B}/M$ is a simple (and continuous) B-module extending the action of A on V and containing V as a dense subset.

(iii) $\Rightarrow$ *(i)*. Let $a \in \tilde{A}$, $\varphi(a)^{-1} \in \tilde{B} - \varphi(\tilde{A})$. We show that *(iii)* is not satisfied. The element a cannot be left invertible in $\tilde{A}$ (otherwise we would have $\varphi(a)^{-1} \in \varphi(\tilde{A})$). Hence $< a >= \tilde{A}a$ is a proper left ideal in $\tilde{A}$. By Zorn's lemma there is some maximal left ideal N in $\tilde{A}$ containing $< a >$. View $V = \tilde{A}/N$ as an A-module. Since N is maximal, V is simple. Note that $a[1] = [a] = 0$, while $[1] \neq 0$. Hence, even if the kernel of φ acts trivially on V (so V factors to an $\varphi(\tilde{A})$-module), V cannot be contained in any B-module since $\varphi(a)$ is invertible in $\tilde{B}$. This proves the contrapositive of *(iii)* $\Rightarrow$ *(i)*. Note that we did not use the assumption that inv$\tilde{B}$ is open or that $\varphi(A)$ is dense for this.**Q.E.D.**

Corollary 1.5. *Let B be a C^*-algebra, and let A be a dense Fréchet subalgebra. Then the following are equivalent.*

(1). *A is closed under the holomorphic functional calculus in B.*

(2). *Every simple (possibly discontinuous) A-module is contained as a dense subspace of an irreducible *-representation of B on a Hilbert space.*

Proof: Condition (1) is equivalent to the spectral invariance of A in B by Lemma 1.2. Condition (2) is equivalent to condition (*iii*) of Theorem 1.4 if B is a C*-algebra by **[6, Corollary 2.9.6(i)]**. **Q.E.D.**

As a point of interest, we remark that if A is a Q-algebra, then every simple A-module V is continuous for a natural topology. Namely choose $v \in V$ and let N be the set of $a \in \tilde{A}$ such that $av = 0$. Then N is closed by maximality and $V \cong \tilde{A}/N$ gives the topology on V.

Example 1.6. For an example where φ is not injective, and A is SI in B, let A be power series in two variables, $\mathbf{C}[[\mathbf{X}, \mathbf{Y}]]$, and let B be power series in one variable, $\mathbf{C}[[\mathbf{X}]]$. Define

$$\varphi(\sum c_{nm} X^n Y^m) = \sum c_{n0} X^n.$$

The units in both A and B are any series with a nonzero constant term, so A is SI in B. Both A and B are local rings (in the algebraic sense that they have a unique maximal ideal), having only one simple module given by evaluating at $X = Y = 0$. If we place the topology of pointwise convergence of coefficients on A and B, then both algebras are Q-algebras, since the set of series with zero constant term is clearly closed.

§2 Spectral Invariance of $M_n(A)$

If A is an algebra, let $M_n(A)$ denote the n by n matrices over A.

Theorem 2.1. *Let $\varphi\colon A \to B$ be an algebra homomorphism with dense image, and assume that B is a Q-algebra. Then A is SI in B iff $M_n(A)$ is SI in $M_n(B)$.*

Proof: We prove the forward direction. Assume that A is SI in B. We will show that condition (*iii*) of Theorem 1.4 holds for the pair $M_n(A)$, $M_n(B)$. We first need a lemma.

Lemma 2.2. *Let A be any algebra, V a simple $M_n(A)$-module. Then V is a direct sum*

$$V = \bigoplus_{1}^{n} V_0$$

where V_0 is a simple A-module.

Proof: Simple exercise using matrix units.**Q.E.D.**

Let V be a simple $M_n(A)$-module. By the lemma, $V = \bigoplus_1^n V_0$ where V_0 is a simple A-module. Since A is SI in B and B is a Q-algebra, we can extend V_0 to a B-module W_0. Then $W = \bigoplus_1^n W_0$ is an $M_n(B)$-module in a natural way, containing V. Hence $M_n(A)$ is SI in $M_n(B)$. **Q.E.D.**

Since $(iii) \Rightarrow (i)$ in Theorem 1.4 did not require B to be a Q-algebra, we did not need to check that $M_n(B)$ is a Q-algebra in Theorem 2.1. It is true by [**12, Lemma 2.1**] that if B is good, then $M_n(B)$ is good. However, I do not know if this is true with "good" replaced by "Q-algebra".

Corollary 2.3. *If A is a dense Fréchet subalgebra of a C^*-algebra B, then Theorem 2.1 is true with "spectral invariant" replaced by "closed under the holomorphic functional calculus".*

Remark 2.4. B. Gramsch has an unpublished proof of Theorem 2.1 which works if B is a good topological algebra with jointly continuous multiplication. This proof was also noticed by Bost in [**3, Proposition A.2.2**], and is very similar to the proof of [**7, Proposition 1.8.4**]. The proof fits roughly into the following two steps. Step 1 was first proven by Swan in [**12, Lemma 1.1**].

Step 1. Assume both A and B are unital (the nonunital case follows easily from the unital case). For simplicity, we assume $n = 2$. Since B is a good topological algebra with jointly continuous multiplication, we can find a neighborhood V of the identity in $M_n(B)$ such that

$$\begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \in V$$

implies b_{11}, b_{22}, and $b_{22} - b_{21}b_{11}^{-1}b_{12}$ are all invertible in B (note we have used the fact that inversion is continuous here). If A is SI in B and

$$[a_{ij}] = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in V \cap M_n(A),$$

then a_{11}, a_{22}, $a_{22} - a_{21}a_{11}^{-1}a_{12}$ are all invertible in A. So

$$\begin{pmatrix} a_{11}^{-1} & 0 \\ 0 & (a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} \end{pmatrix} \begin{pmatrix} 1 & -a_{12}(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -a_{21}a_{11}^{-1} & 1 \end{pmatrix}$$

is easily seen to be an inverse for $[a_{ij}]$. (To construct this inverse, we have simply done elementary row and column operations to put the matrix into diagonal form, then cleared the diagonal.)

Step 2. Next we note that if we can find such a neighborhood V as in step 1, then $M_n(A)$ is SI in $M_n(B)$. For let $a \in M_n(A) \cap \mathrm{inv} M_n(B)$. By density, there are $a_1, a_2 \in M_n(A)$ such that $a_1 a \in V$, $a a_2 \in V$. It follows that a is invertible in $M_n(A)$. This is the argument for the last statement of [**8, Lemma 5.3**], which is also given in [**3, Proposition A.2.2**].

Question 2.5. Do there exist Q-algebras without continuous inversion ? The field without continuous inversion described in [**13, IX.2**] may be such an example, however I have not been able to fill in the details of this example, and have heard that others have had the same problem.

§3 Examples

Example 3.1. We give an example of a non SI dense Banach subalgebra of a C*-algebra with an open group of invertible elements.

Let $B = C[-1, 1]$. Let A be the set of elements of B which extend to analytic functions on the open disc $\{ z \in \mathbf{C} \mid |\mathbf{z}| < \mathbf{1} \}$, continuous on the boundary. The sup norm over the whole disc makes A a Banach algebra. The subalgebra A is not SI since $i + z$ is invertible in B. For a simple A-module not extending to any B-module, let $V = \mathbf{C}$, $fc = f(-i)c$.

We use Theorem 1.4 to show that some dense subalgebras are *not* spectral invariant.

Example 3.2. Let $\mathbf{Z}$ act on $\mathbf{T}$ by an irrational rotation α. Let B be the C*-crossed product $C^*(\mathbf{Z}, \mathbf{C}(\mathbf{T}), \alpha)$, and let $A = C_c^\infty(\mathbf{Z} \times \mathbf{T})$, the convolution algebra of C^∞ functions with compact support on $\mathbf{Z} \times \mathbf{T}$. The representation of A on $C^\infty(\mathbf{T})$ given by

$$(Ff)(z) = \sum_{n \in \mathbf{Z}} F(n, z) e^{-n} f(e^{-\iota \alpha n} z), \tag{3.3}$$

where $F \in A$, $f \in C^\infty(\mathbf{T})$ and $z \in \mathbf{T}$, is not contained in any B-module, since the character $n \mapsto e^{-n}$ does not give a character of $C^*(\mathbf{Z})$. We show that (3.3) defines an algebraically irreducible representation of A. If $f \in C^\infty(\mathbf{T})$ and f is not identically zero, then $|f| \in C^\infty(\mathbf{T})\mathbf{f} \subseteq \mathbf{Af}$. So Af contains a nonnegative and nonzero function, and all of its translates by multiples of α. Since $\mathbf{T}$ is compact, the sum of finitely many of these translates never vanishes on $\mathbf{T}$. Call this sum g. Then $C^\infty(\mathbf{T}) \subseteq \mathbf{C}^\infty(\mathbf{T})\mathbf{g} \subseteq \mathbf{Af}$. Thus the representation is algebraically irreducible. By Theorem 1.4, A is not spectral invariant in B. We remark that if the action of $\mathbf{Z}$ on a manifold M is *proper*, then the convolution algebra $C_c^\infty(\mathbf{Z} \times \mathbf{M})$ *is* SI in the C*-crossed product [**1, Appendix**].

Example 3.4. Let G be the real $ax+b$ group, defined as $\mathbf{R}^2$ with multiplication $(a,b)(c,d) = (a+c, e^a d + b)$. We define a dense subalgebra of $C^*(G)$ consisting of "exponentially decreasing" Schwartz functions, which we show is not spectral invariant. Define $\omega(a,b) = 1+e^{|a|}+|b|+|e^{-a}b|$. Then ω is a submultiplicative function (or weight) on G, and we define the ω-rapidly vanishing differentiable functions on G to be those differentiable functions ψ from G to $\mathbf{C}$ satisfying

$$\int_G \omega^d(g)\, |D\psi(g)|\, dg < \infty \tag{3.5}$$

for each differential operator D on G and each natural number d. Here dg denotes the left Haar measure on G. This space $S^\omega(G)$ is then a dense m-convex Fréchet *-subalgebra of $C^*(G)$ [**11, §1**]. Left Haar measure on G is given by $d\mu(a,b) = e^{-a}dadb$, where $(a,b) \in \mathbf{R^2}$ and $dadb$ is Lebesgue measure on $\mathbf{R^2}$. The character

$$\psi \mapsto \int_{\mathbf{R^2}} \psi(a,b)dadb = \int e^a \psi(a,b) d\mu(a,b) \tag{3.6}$$

defines a simple $S^\omega(G)$-module not contained in any $C^*(G)$-module, since the character $(a,b) \mapsto e^a$ of G is not unitary. So $S^\omega(G)$ is not SI in $C^*(G)$.

We conclude with two examples of what happens in Theorem 1.4 (*i*) ⇒ (*iii*) when the condition that B is a Q-algebra is dropped.

Example 3.7. We give an example of a pair of topological algebras such that A is SI and dense in B, $\mathrm{inv}\tilde{B}$ is not open, and there is a simple A-module not contained in any B-module. Let $B = \mathbf{C}[\mathbf{X},\mathbf{Y}]$, $A = \mathbf{C}[\mathbf{Z},\mathbf{Y}]$, the sets of complex polynomials in X, Y and Z, Y respectively. Identify A with a subalgebra of B by mapping Z to $X(1-Y)$.

We give B the following locally convex Hausdorff topology. Let $\mathbf{C}[[\mathbf{X},\mathbf{Y}]]$ be the algebra of power series in two variables. If $n, m \in \mathbf{N}$, we define ideals $I_{n,m} =< X^n, Y^m >$

in $\mathbf{C}[[\mathbf{X}, \mathbf{Y}]]$. Then we say that $f_k \to f_0$ in $\mathbf{C}[[\mathbf{X}, \mathbf{Y}]]$ if and only if $f_k - f_0$ is eventually in every $I_{n,m}$. Let $B \subseteq \mathbf{C}[[\mathbf{X}, \mathbf{Y}]]$ have the relative topology. With this topology on B, we see that $X(1-Y)(1+Y+\dots Y^n)$ converges to X as $n \to \infty$, so $X \in \overline{A}^B$ and A is dense in B. The units in both algebras are the constants, so A is SI in B, and inv$\tilde{B}$ is not open. Evaluation at Z, $Y = 1$ gives a simple A-module contained in no B-module.

Example 3.8. We investigate further what happens in Theorem 1.4 if we drop the hypothesis that inv$\tilde{B}$ is open. It is still true (in the implication $(i) \Rightarrow (iii)$) that if an extension exists it can be taken simple. However, we show by example that extensions may always exist, but we may not be able to take our extension to be continuous, or have dense inclusion. In fact, in our example V is one-dimensional, but any extension W is discontinuous and infinite dimensional.

1. Let $B = s(\mathbf{N})$, the sequences of (possibly unbounded) complex numbers with pointwise multiplication and the topology of pointwise convergence. This space is a complete, unital, locally convex, m-convex Fréchet algebra over $\mathbf{C}$ (see **[9, Example 3.7]**). An element of B is invertible iff it is never zero. One easily finds a sequence of non-invertible elements converging to 1, so inv$B$ is not open. Hence we know by **[14]** that B will have discontinuous infinite dimensional simple modules.

2. We characterize all simple B-modules, or equivalently all maximal ideals. Let U denote the set of ultrafilters on $\mathbf{N}$ (see **[4]**). For any $F \in U$ we define an ideal in B by

$$I_F = \{\, b \in B \mid \quad \exists S \in F(b_n = 0 \Longleftrightarrow n \in S) \quad \} \tag{3.9}$$

One easily checks that this is a maximal ideal, and that every maximal ideal arises in this way. For each $p \in \mathbf{N}$, we have a maximal ideal $I_p = \{\, b \in B \mid b_p = 0 \,\}$. Such a maximal ideal corresponds to the principal ultrafilter $F_p = \{\, S \subseteq \mathbf{N} \mid \mathbf{p} \in \mathbf{S} \,\}$. There are many non-principal ultrafilters on $\mathbf{N}$ (see **[4, Theorem 7.1]**).

Any non-principal ultrafilter F contains all cofinite sets so that I_F contains all finitely supported sequences in B. Thus I_F is a dense maximal ideal in B and B/I_F is a discontinuous simple B-module. One can find elements of B/I_F which are not scalar multiples of the identity, so that B/I_F is a proper field extension of the algebraically closed field $\mathbf{C}$. Hence B/I_F is infinite dimensional over $\mathbf{C}$.

3. Now we define our dense subalgebra A to be the algebra of all finitely supported sequences $s_f(\mathbf{N})$ with unit adjoined. Let V be the trivial A-module, defined by $V = \mathbf{C}$

with action $(s+\lambda)z = \lambda z$, for $s \in s_f(\mathbf{N})$, $\lambda \in \mathbf{C}$, and $z \in V$. Clearly $s_f(\mathbf{N})$ is the kernel of this representation, so the kernel of any irreducible representation of B containing V is a maximal ideal containing $s_f(\mathbf{N})$. Hence any extension of V to a simple B-module W must be of the form $W = B/I_F$ for some non-principal ultrafilter F. Hence there is no extension of V to a continuous simple B-module.

4. We note, however, that every simple A-module does have an extension to a B-module (so that $(i) \Rightarrow (iii)$ does not fail). To show this, we classify all simple A-modules by noting that $s_f(\mathbf{N})$ is dense and SI in the C*-algebra $c_0(\mathbf{N})$ of sequences tending to zero at infinity with the sup norm topology. By Theorem 1.4, all simple A-modules extend to simple $\tilde{c}_0(\mathbf{N})$- modules. Thus the only simple A-modules are the point evaluations and the trivial module V defined in **3**, all of which extend to B-modules.

References

[1] P. Baum, A. Connes, *Chern Character for Discrete Groups*, Collection: A fete of topology, Academic press, Orlando, Florida, 1988, pp. 163-232.

[2] B. Blackadar, K-theory for operator algebras, Springer-Verlag, New York, 1986.

[3] J.B. Bost, *Principe D'Oka, K-Theorie et Systems Dynamiques Non-commutative*, Invent. Math, **101** (1990), 261-333.

[4] W.W. Comfort, S. Negrepontis, The Thoery of Ultrafilters, Springer-Verlag, Berlin/Heidleberg, 1974, pp.142-144.

[5] A. Connes, *An Analogue of the Thom Isomophism for Crossed Products of a C*-algebra by an Action of* $\mathbf{R}$, Adv. in Math. **39** (1981), 31–55.

[6] J. Dixmier, C* algebras, North-Holland Publishing Co., Amsterdam New York Oxford,1982.

[7] T. Palmer, Banach Algebras and the General Theory of *-Algebras, Cambridge University Press Encyclopedia of Mathematics Series, in preparation.

[8] B. Gramsch, *Relative Inversion in der Störungstheorie von Operatoren und* Ψ-*Algebren*, Math Ann **209** (1984), 27–71.

[9] E. Micheal, *Locally multiplicatively convex topological algebras*, Mem. Amer. Math. Soc., **11** (1952).

[10] N.C. Phillips, *K-theory for Fréchet algebras*, Intl. Jour. Math. **2 (1)** (1991), 77-129.

[11] L. Schweitzer, *Dense m-convex Fréchet Subalgebras of Operator Crossed Products by Lie Groups*, preliminary version (1990).

[12] R. G. Swan, *Topological Examples of Projective Modules*, Trans. Amer. Math. Soc.**230** (1977), 201-234.

[13] L. Waelbroeck, Topological Vector Spaces and Algebras, Springer-Verlag, Berlin Heidelberg New York, 1971.

[14] W. Zelazko, *On maximal ideals in commutative m-convex algebras*, Studia Math **T.LVIII** (1976), 291-298.